

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Journal of Otorhinology and Allied Science

Journal homepage: <https://www.joas.co.in/>

Case Report

Absent stapedius muscle, tendon and pyramid – An anomaly revisited

Abhishek Bhardwaj¹, Rachit Sood^{1,*}, Manu Malhotra², Madhu Priya¹,
Nivedhan Ravichandran¹, Akhilesh Chandra Yadav¹, Ramesh Prasath¹, Jay Dave¹

¹Dept. of Otorhinology and Allied Science, All India Institute of Medical Science Rishikesh, Uttarakhand, India

²Dept. of Otorhinology (ENT), All India Institute of Medical Science Rishikesh, Uttarakhand, India



ARTICLE INFO

Article history:

Received 14-12-2022

Accepted 21-12-2022

Available online 04-01-2023

Keywords:

Stapedius muscle

Tendon

Pyramid

Intraoperative

ABSTRACT

Congenital absence of stapedius is an exquisite entity, with an incidence of 0.5% and very few cases reported in live patients. We herein report a case of a middle-aged female presented with bilateral ear discharge and decreased hearing for more than a year, which, on examination, showed a large central perforation bilaterally, and an audiogram preoperatively revealed an average hearing loss of 45 dB and air-bone gap of 35dB bilaterally. Right endoscopic type I tympanoplasty was done, which unveiled absent stapedius muscle, tendon, and pyramid on the right side. The absent stapedius should circumspect the operating surgeon to look for other associated anomalies such as tympanosclerosis, fixed footplate and aberrant course of the facial nerve.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Congenital middle ear anomalies are uncommon (1 in 10,000 live births). The stapedius muscle contributes significantly to acoustic reflex in response to loud sounds and protects the inner ear from acoustic trauma. Various anomalies of the stapedius unit have been reported, such as double or ectopic stapedius, ossified tendon, or absence of stapedius. Congenital absence of stapedius is an *extremely rare entity*, with an *incidence of 0.5%* and very few cases reported in live patients. These are often accompanied by other middle ear anomalies, such as congenital aural atresia, aberrant facial nerve or dysplastic ossicular chain, fixed footplate or tympanosclerosis.¹⁻⁵

2. Discussion

A middle-aged female presented with bilateral ear discharge for two years and decreased hearing in both ears for




Fig. 1: Absent pyramid, stapedius muscle and tendon in the right ear (1 - tympanic segment of facial nerve, 2 – handle of malleus, 3 – long process of incus, 4 – round window, 5 – posterior crura of stapes)

* Corresponding author.

E-mail address: nivedhanmbbs@gmail.com (R. Sood).




Fig. 2: Normal pyramid, stapedius muscle and tendon in the left ear (1 - tympanic segment of facial nerve, 2 - handle of malleus, 3 - long process of incus, 4 - round window, 5 - posterior crura of stapes, 6 - stapes footplate, 7 - pyramid, 8 - stapedius muscles and tendon)

one year. On examination, a large central perforation was present bilaterally. The audiogram revealed bilateral moderate conductive hearing loss with an average hearing loss of 45 dB and an air-bone gap of 35 dB. Right endoscopic type I tympanoplasty was done. The stapedius muscle, tendon, and pyramid were absent on the right side and present on the left side. (Figures 1 and 2) The round window reflex was present on the right side. The patient had no complaints of hyperacusis. The graft healed well six weeks postoperatively, and hearing improved.

The case reported by Magnuson et al. had an absent stapedius with a well-developed pyramid but an aberrant facial nerve and absent footplate, and oval window. However, in our patient, it was unilateral, asymptomatic, and without any other anomalies. In his case series of patients with absent stapedius tendon, Nassiri et al. (2021) found congenital stapes footplate fixation in all eight patients.^{6,7} However, this was not seen in our case, as the footplate was mobile.

Stapes is the first ossicle to appear and is first seen at around the fifth week of gestation. Conventionally, it was believed that stapes have a dual origin, with its suprastructure originating from the second arch and footplate from the otic capsule. However, this theory has been rejected, and stapes is known to originate from the second arch. The posterior belly of the digastric muscle gives off fascicles which pass through the stylomastoid foramen into the tympanic cavity and gives stapedius muscles through the neck of stapes. This also explains the similar innervation of the digastric and stapedius posterior belly and their common origin from the second pharyngeal arch.^{4,5,8,9}

Preoperative identification of this entity is challenging even with High-Resolution Computed Tomography. Due to this and its rare incidence, the diagnosis of absent stapedius is almost always made *intraoperatively*. The pyramid plays a vital role and gives bony projections, i.e. the chordal ridge and the ponticulus. These landmarks divide the superior retrotympaanum into facial recess, posterior tympanic sinus, lateral tympanic sinus and sinus tympani. The absence of the pyramid can lead to the distorted anatomy of these recesses. Thorough knowledge and evaluation of this should be done preoperatively on radiology by the operating surgeon as these are the most common sites for residual disease.

3. Results and Conclusion

1. Absent stapedius muscle is a rare entity with an incidence of 0.5%.
2. This is usually associated with other middle ear anomalies and can present with hyperacusis and conductive hearing loss. However, in our case, it was unilateral, asymptomatic, and isolated (without any other anomalies).
3. The absent stapedius should alert the operating surgeon to look for other associated anomalies such as tympanosclerosis, fixed footplate and aberrant course of the facial nerve.

4. Source of Funding

None.

5. Conflict of Interest

None.


References

1. Schwager K. Reconstruction of middle ear malformations. *GMS Curr Top.* 2007;6:1–16.
2. Philippon D, Laflamme N, Leboulanger N. Hearing Outcomes Functional Surgery for Middle ear Malformations. *Otol Neurotol Publ Am Otol Soc Am Neurotol Soc Eur Acad Otol Neurotol.* 2013;34(8):1417–20.
3. Dey S, Dutta M, Misra S. Congenital absence of stapedius muscle and tendon: Rare finding in two cases. *Indian J Oncol.* 2017;23(1):43–5.
4. Dalmia D, Behera S. Congenital Absence Of Stapedius Muscle And Tendon: Case Report With Review Of Literature. Author: Samyadipta Dey, Mainak Dutta, Saibal Misra. *J Clin Diagn Res.* 2022;doi:10.7860/jcdr/2022/49726.15820.
5. Kanona H, Virk JS, Kumar G, Chawda S, Khalil S. A Rare Stapes Abnormality. *A Rare Stapes Abnormality.* 2015;p. 387642. doi:10.1155/2015/387642.
6. Magnuson T, Har-El G. Middle ear anomalies. *Otolaryngol Head Neck Surg Off J Am.* 1994;111(6):853–4.

7. Nassiri AM, Benson JC, Doerfer KW, Perkins EL, Sweeney AD, Patel NS. Absent pyramidal eminence and stapedial tendon associated with congenital stapes footplate fixation: Intraoperative and radiographic findings. *Am J Otolaryngol*. 2021;42(6):103144. doi:10.1016/j.amjoto.2021.103144.
8. Vázquez JF. Development of the stapes and associated structures in human embryos. *J Anat*. 2005;207(2):65–73. doi:10.1111/j.1469-7580.2005.00441.x.
9. Hough J. Congenital Malformations of the Middle Ear. *Arch Otolaryngol*. 1985;p. 1–9. Available from: <https://www.ajronline.org/doi/pdfplus/10.2214/ajr.144.3.501>.


Author biography

Abhishek Bhardwaj, Associate Professor

Rachit Sood, Senior Resident  <https://orcid.org/0000-0002-4644-1296>

Manu Malhotra, Professor

Madhu Priya, Additional Professor

Nivedhan Ravichandran, Junior Resident  <https://orcid.org/0000-0002-3785-9658>

Akhilesh Chandra Yadav, Junior Resident

Ramesh Prasath, Junior Resident

Jay Dave, Junior Resident

Cite this article: Bhardwaj A, Sood R, Malhotra M, Priya M, Ravichandran N, Yadav AC, Prasath R, Dave J. Absent stapedius muscle, tendon and pyramid – An anomaly revisited. *IP J Otorhinolaryngol Allied Sci* 2022;5(4):138-140.